


Demographic snapshot

	San Jose	SCC
Population size	945,942	1,781,642
<i>Race/ethnicity</i>		
African American	3%	2%
Asian/Pacific Islander	32%	32%
Latino	33%	27%
White	29%	35%
Foreign-born	39%	37%
Speaks a language other than English at home	56%	52%
Single parent households	8%	7%
Households with children	41%	39%
Average household size	3.09	2.90
<i>Age</i>		
0-5 years	9%	8%
6-11 years	8%	8%
12-17 years	8%	8%
18-24 years	10%	9%
25-34 years	15%	15%
35-44 years	16%	16%
45-54 years	14%	15%
55-64 years	10%	10%
Ages 65 and older	10%	11%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: U.S. Census Bureau, 2010 Census and 2010-2014 American Community Survey 5-Year Estimates



Economic and educational opportunities

Income and job opportunities

	San Jose	SCC
Median household income	\$83,787	\$93,854
Unemployed	4.2%	3.7%
<i>Poverty</i>		
Families below 185% FPL	20%	16%
Children (ages 0-17) below 185% FPL	31%	25%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: U.S. Census Bureau, 2010 Census and 2010-2014 American Community Survey 5-Year Estimates; State of California Employment Development Department, December 2015 downloaded 2/26/16

Opportunities for high quality and accessible education

	San Jose	SCC
Children (ages 3-4) enrolled in preschool or nursery school	52%	57%
Educational attainment (ages ≥ 25 years)		
Less than high school	17%	13%
High school graduate	18%	15%
Some college or associates degree	26%	24%
College graduate or higher	38%	47%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: U.S. Census Bureau, 2010-2014 American Community Survey 5-Year Estimates

Healthy and safe environment

Safe, sustainable, accessible, and affordable transportation options

	San Jose	SCC
Number of vehicle-pedestrian injury collisions, 1 year	300	478
Number of vehicle-bicycle injury collisions, 1 year	330	636
Number of motor vehicle collisions, 1 year	3599	6669
Population that resides within a ½ mile of a transit stop with a headway of 15 minutes or less during peak commute hours	72%	65%
Residents who commute to work		
Drove alone	78%	76%
Carpooled	11%	10%
Public transportation	4%	4%
Other	8%	10%
Travel time to work over 30 minutes	40%	37%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: U.S. Census Bureau, 2010 Census and 2010-2014 American Community Survey 5-Year Estimates; Statewide Integrated Traffic Records System, 2004-2013, October 5, 2015; 2012 Transit Stops from the Metropolitan Transportation Commission

Affordable, accessible, and nutritious foods

	San Jose	SCC
Households receiving CalFresh benefits	7%	5%
Average distance (miles) to nearest full-service grocery store	0.53	0.56
Average distance (miles) to nearest farmers' market	1.35	1.60
Number of fast food outlets per square mile	2.8	2.8
Adults who ate 2+ servings of fruit per day in the past 30 days	26%	27%
Adults who ate 3+ servings of vegetables per day in the past 30 days	18%	19%
Adults who ate fast food at least weekly in the past 30 days	42%	38%
Adults whose household shops for fruits and vegetables within community/neighborhood	90%	92%
Adult(s) in family ever received food from a church, a food pantry, or food bank in the past 12 months	14%	11%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: Santa Clara County Information Services Department Farmers Markets 2016; InfoUSA, April 2014; Santa Clara County Public Health Department, 2013-14 Behavioral Risk Factor Survey

Affordable and high quality housing

	San Jose	SCC
Households occupied by renters	43%	43%
Households with gross rent 30% or more of household income	52%	46%
Overcrowded households	10%	8%
Multi-unit housing	31%	33%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: U.S. Census Bureau, 2010-2014 American Community Survey 5-Year Estimates

Access to affordable and safe opportunities for physical activity

	San Jose	SCC
Adults who met CDC recommendations for aerobic physical activity in the past month	53%	58%
Average distance (miles) to the nearest park or open space	0.29	0.29

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

The Centers for Disease Control and Prevention (CDC) suggests that adults ages 18 and older get at least 2 hours and 30 minutes (150 minutes) of moderate intensity aerobic activity (i.e., brisk walking) every week for good health; 1 hour and 15 minutes (75 minutes) of vigorous intensity aerobic activity (i.e., jogging or running); or an equivalent mix of moderate and vigorous intensity activity.

Sources: Santa Clara County Parks and Recreation, 2016; Santa Clara County Public Health Department, 2013-14 Behavioral Risk Factor Survey

Tobacco and smoke-free

	San Jose	SCC
Adults who are current smokers	12%	10%
Adults who smelled tobacco smoke drifting into home from nearby apartments or from outside in the past 7 days (among adults living in multi-unit housing)	30+*%	28%
Number of tobacco retail outlets per square mile	3.9	3.5

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: Santa Clara County Public Health Department, 2013 Behavioral Risk Factor Survey; California State Board of Equalization, April 2016

Safe communities, free of crime and violence

	San Jose	SCC
Average number of violent crimes within 1 mile	21.8	16.04
Number of alcohol retail outlets per square mile	2.7	2.7
Adults who report neighborhood crime, violence, and drug activity is somewhat of or a major problem	54%	42%
Adults with any firearms now kept in or around home	10%	11%
Adults who were ever hit, slapped, pushed, kicked, or hurt in any way by an intimate partner	12%	12%

Notes: * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Source: Public Engines, Inc., January 1, 2012 to December 31, 2012; California Department of Alcoholic Beverage Control, off site alcohol retailers, April, 2016; Santa Clara County Public Health Department, 2013-14 Behavioral Risk Factor Survey

Health status

	San Jose	SCC
<i>Maternal, child, and infant health</i>		
Births per 1,000 people	14.5	13.5
Low birth weight infants	7%	7%
Preterm births	9%	9%
Overweight or obese mothers in first trimester of pregnancy	42%	38%
Mothers who received early and adequate prenatal care	73%	74%
Teen live births per 1,000 females, ages 15-19	25.3	19.2
<i>Health behaviors</i>		
Adults with health coverage (ages 18-64)	84%	85%
Adults with dental insurance	62%	64%
Adults with fair or poor self-rated health	22%	19%
Adults who have health condition(s) that limit(s) activities	20%	19%
Adults who experienced frequent mental distress in the past 30 days	11%	9%
Adults who were ever diagnosed with depression	15%	14%
Adults who are obese	22%	20%
Adults who received a flu shot or nasal flu vaccine in the past 12 months	41%	42%
Adults who engaged in binge drinking in the past 30 days	15%	14%
Adults who were ever diagnosed with high blood pressure (hypertension)	28%	27%
Adults who were ever diagnosed with asthma	14%	14%
Adults who were ever diagnosed with diabetes	7%	8%
Adults who had 1+ falls in the past 3 months (ages 45+)	14%	15%
<i>Mortality</i>		
Life expectancy	82.0	83.4
Cancer deaths per 100,000 people	150.4	140.3
Heart disease deaths per 100,000 people	126.2	118.8
Alzheimer's disease deaths per 100,000 people	34.1	34.6
Stroke deaths per 100,000 people	30.5	27.8
Chronic lower respiratory disease deaths per 100,000 people	29.0	25.5
Unintentional injury deaths per 100,000 people	26.4	23.4
Diabetes deaths per 100,000 people	29.2	23.3
Influenza and pneumonia deaths per 100,000 people	15.9	14.3
Hypertension deaths per 100,000 people	18.3	14.6

Notes: Frequent mental distress is an indicator of general mental health and is defined as 14 or more days during the past 30 days on which mental health was not good. Early and adequate prenatal care, also known as the Kotelchuck Index is defined as care initiated by the fourth month and receiving at least 80% of recommended prenatal visits. Chronic lower respiratory disease includes asthma, COPD, bronchitis, and emphysema. * + indicates estimate is statistically unstable due to a relative standard error of greater than 30% (*) or less than 50 respondents in the denominator (+). These estimates should be viewed with caution and may not be appropriate to use for planning or policy purposes.

Sources: Santa Clara County Public Health Department, 2009-2013 Birth Statistical Master File; Santa Clara County Public Health Department, 2013 Behavioral Risk Factor Survey; Santa Clara County Public Health Department, 2009-2013 Death Statistical Master File; U.S. Census Bureau, 2010 Census; Summary File 1, Table PCT12.